

Effect of Different Mining Dust on the Vegetation of District Balaghat, M.P - A Critical Review

Jyotish Katare¹, Mohnish Pichhode², Kumar Nikhil³

^{1,2}Department of Biotechnology, Sardar Patel College of Technology, Balaghat, Rani Durgavati University Jabalpur, Madhya Pradesh, India

³Principal Scientist, EMG, CSIR- Central Institute of Mining and Fuel Research (CIMFR), Dhanbad, Jharkhand, India

Abstract: *Industrialization had increased requirement of more energy with mining for more minerals resultant pollution. Soil pollution due to different mining dust has become problem in district Balaghat, M.P beside that dust deposition on the surface of vegetation affects photosynthesis and growth of common plants nearby the mining areas. This review articles describes briefly the various mining sites in Balaghat district of Madhya Pradesh, India and dust chemical characteristics which had many effects on vegetation and soil. The physical and chemical characteristics of a range of mining dust types are explained and its effects on photosynthesis, respiration and transpiration. Plants growing on this atmosphere show a reduction in growth performance and yield. Visible injury symptoms and decreased in productivity on vegetation is well noticed. Most of the plant community structure is altered. However, there have been very few detailed studies on natural and semi-natural systems and some dust types are also much understudied. Recommendations for future research are made in order to overcome this knowledge gap.*

Keywords: Dust, Growth performance of Vegetation, Mining in district Balaghat, M.P, Soil pollution.

1. Introduction

This paper briefly explains the effects of mining dust pollution on plant communities. Dusts consist of solid matter in a minute and fine state of subdivision so that the particles are small enough to be raised and carried by wind. They may originate from many sources. A large number of mines and mining industrial process can produce particulate emission (Fennelly, 1975). The main processes that regularly cause problems, however, are those concerned with mineral extraction. This range from the quarrying itself to the various processing operations e.g. copper, manganese, limestone, marble and aluminum ore processing with cement manufacturing. Mining dust suppression in this operation is more difficult and dust levels can be very high. Mineral extraction is increasing in many countries in order to meet increased construction demand (Harris, 1975, Stanton 1989). Industrialization, urbanization, economic growth and associated increase in metal and minerals demands have resulted in a profound deterioration of air quality in developing countries like India.

Although minerals and heavy metals are naturally present in the soil, geologic and anthropogenic activities increase the concentration of these elements to amounts that are harmful to plants. Some of these activities include mining smelting of metals, processing of minerals also. Growth reduction as a result of changes in physiological and biochemical processes in plants growing on mining polluted soils has been recorded (Chatterjee & Chatterjee, 2000).

This paper shows a comprehensive review of literatures available on the effect of mining dust on plants with their communities and soil. Initially the characteristics of various dust types are described, highlighting those factors that are important in determining the likely impact its deposition may have. The effect of mining dust is considered according to vegetation type followed by a discussion of the effects of

mining dust on plant species, crop species and on a range of natural and semi-natural vegetation types.

Further, the objective of this study is to know the influence of mining activity on the concentration of dust in air, soil and vegetation surrounding mining sites at Balaghat district, Madhya Pradesh, India.

2. Detail of sites

2.1 About different mining sites in district Balaghat, M.P.

In Madhya Pradesh, Balaghat district has a huge mineral deposits and also prosperous with forest. Balaghat is located in the southern part of Jabalpur division. It occupies the south eastern portion of the satpura range and the upper valley of the Wainganga River. The district extends from 21°19' to 22°24' north latitude and 79°31' to 81°3' east longitude. The total area of the district is 9,245km² (Figure.2.1 and 2.2).

About 80% of the manganese production in India comes from Balaghat district by MOIL, and copper deposit at Malanjkhand, Balaghat is regarded 70% as the largest in the country. In fact it is largest open-cast mining in Asia. Bauxite, Kainite, Marble, Dolomite, Clay and Limestone are the other main minerals of the Balaghat district (Table.2.1).

The lush green forest of Balaghat district and Kannah National park is also important as mining of different minerals present in and around.

Table 2.1: Mineral Resources of District Balaghat, M.P, India

S. N.	Minerals	Village	Tehsil	Uses
01.	Copper ore (Chalcopyrite, Pyrite, Cuprite)	Malanjkhanda	Baihar	Used in wire, plates etc.,
02.	Manganese ore (Pyrolusite, Bryonite, Magnite)	Ukwa, Tirodi, Hirapur, Bharweli, Manegaon, Miragpur	Baihar Katangi Balaghat	Used in steel and alloys.
03.	Cement Raw	Duglai, Hatri, Kandai, Deodongla, Jangla, Piprahi	Baihar	Used for Cement
04.	Marble Dolomite	Bhagatpur latri, Sitapchoro, Dulhapur	Balaghat Baihar Katangi	Used in Tiles
05.	Bauxite (Alumina)	Mardi Dadar, Toursi Dadar, Garhi Dadar	Baihar	used in Refractory
06.	Limestone	Tirodi	Katangi	Used in Cement.

3. Source of Contamination

There are different sources of dust in the environment such as (1) Natural source (2) Agriculture source (3) Industrial Source (4) Atmospheric source (5) Mining Source etc. Dust can be emitted into the environment by both natural and anthropogenic activities. The major causes of emission are the anthropogenic sources specifically mining operations (Nriagu 1989). In some cases, even long after the mining activities have ceased the emitted metals continue to persist in the environment. Peplow (1999) reported that hard rock mine operate from 5-15 years until the minerals are depleted, but metal contamination that occurs as a consequence of hard rock mining persist for hundred of year after the cessation of mining operations. Heavy metal and minerals are emitted both in elemental and in compound (organic and inorganic) forms. Anthropogenic sources of emission are the various industrial point sources including former and present mining site.

4. Effect of Mining Dust on Plants

Plants are often sensitive both to deficiency and to the excess availability of some heavy metal ion as essential micronutrient, while the same at higher concentrations and ions are strongly poisonous to the metabolic activities. Research shown significant adverse effects of toxic mining dust on plants (Reeves and Baker 2000, Fernandes and Henriques 1991) and agricultural soil has become a critical environmental concern due to their potential adverse ecological effects. Further, such toxic elements are considered as soil pollutants and their acute and chronic toxic effect on plants grown in such soils.

4.1 Effect of Copper Dust on Plants

Copper (Cu) is considered as a micronutrient for plants (Thomas et al., 1998) and plays important role in CO₂ assimilation and ATP synthesis. Cu is also an essential component of various proteins like plastocyanin of photosynthetic system and cytochrome oxidase of respiratory electron transport chain (Demirevska-kepova et al., 2004). But enhanced industrial and mining activities have contributed to the increasing occurrence of Cu in ecosystems. Cu is also added to soils from different human activities including mining and smelting of Cu-containing ores. Mining activities generate a large amount of waste rocks and tailings, which get deposited at the surface. Excess of Cu in soil plays a cytotoxic role, induces stress and ROS (Stadtman and Oliver 1991). Oxidative stress causes disturbance of metabolic pathways and damage to macromolecules (Hegedus et al., 2001). Copper dust had adverse effect on photosynthesis pigmentations of vegetation around the mining site (Pichhode & Nikhil, 2015 a) with soil and vegetation cumulatively on certain vegetation of district Balaghat, M.P (Pichhode & Nikhil, 2015 b). Excess application of pesticides had resultant in the rise of Cu in soil (Nikki, 2000).



Figure 2.1: Location Map of district Balaghat, M.P, India




Figure 2.2: Geographical location of different mining site in district Balaghat, M.P, India

4.2 Effect of Manganese Dust on Plants

Accumulation of excessive manganese (Mn) in leaves causes a reduction of photosynthetic rate (Kitao et al., 1997). Mn is readily transported from the root to shoot through phloem to other organs after reaching the leaves (Loneragan 1988). Necrotic brown spotting on leaves, petioles and stems is a common symptom of Mn toxicity (Wu 1994). This spotting starts on the lower leaves and progresses with time toward the upper leaves (Horiguchi 1988). With time, the speckles can increase in both number and size resulting in necrotic lesions, leaf browning and death (Elamin and Wilcox 1986). General leaf bronzing and shortening of internodes has been documented in *Cucumis sativus* (cucumber) (Crawford et al., 1989, a and b). Another common symptom is known as "crinkle-leaf", and it occurs in the youngest leaf, stem and petiole tissue. It is also associated with chlorosis and browning of the tissue (Wu 1994; Bachman and Miller 1995). Roots exhibiting Mn toxicity are commonly brown in colour (Le Bot et al., 1990; Foy et al., 1995) and sometimes crack (Foy et al., 1995). Chlorosis in younger leaves by Mn toxicity is thought to be caused through Mn-induced Fe deficiency (Horst 1988). Excess Mn is reported to inhibit synthesis of chlorophyll by blocking a Fe-concerning process (Clarimont et al., 1986). Manganese toxicity in some species starts with chlorosis of older leaves moving toward the younger leaves with time (Bachman and Miller 1995). This symptom starts at the leaf margins progressing to the interveinal areas and if the toxicity is acute, the symptom progresses to marginal and interveinal necrosis of leaves (Bachman and Miller 1995).

4.3 Effect of Iron on Plants

Iron as an essential element for all plants has many important biological roles in the processes as diverse as photosynthesis, chloroplast development and chlorophyll biosynthesis. Iron is a major constituents of the cell redox system such as heme proteins including cytochromes, catalase, peroxidase and leghemoglobin and iron sulfur proteins including ferredoxin, aconitase and superoxide dismutase (SOD) (Marschner 1995).

4.4 Effect of Cement Dust on Plants

The cement industries also plays a vital role in the imbalances of the environment and produces air pollution hazards (Stern 1976). These dust particulates are causing large scale deforestation destruction of biota (Panda, et al., 1996) and other natural resources. Among these deposition of cement kiln dust in large quantities around cement factories causes changes in soil's physical and chemical properties (Asubiojo, et al., 1991; Saralabai 1993). The effect of such deposition affects the growth and biochemical characteristics of field crops has also been widely studied (Prasad and Inamadar 1990; Prasad et al., 1991). According to Farmer (1990) cement industrial region are confronted with the problems of alkalization due to high deposition of alkaline cement dust and their ash in the pollution complex. In addition, the growth of quarrying and open-cast mining suggests the deposition on vegetation may be increasing.

4.5 Effect of Bauxite Dust on Plants

Bauxite mining is one such major open cast mining activity which has significant negative impact on the local environment. The major threats of this activity are dust pollution, vegetation loss, forest fragmentation and biodiversity loss, negative impact on water resources, generation of wastelands and social impact. The study revealed that the legal and illegal mining activity has initiated serious environmental degradation in the region. Though mining initially provided job opportunities for limited inhabitants and generated revenue to Government, it would last only for a short period. However, the damage caused to the local ecology as a result of the changed land use is permanent (Rohan and Samant, 2012).

4.6 Effect of Marble Dust on Plants

The paper describes the effect of marble dust on plants. Trees species growing were selected and various morphological characteristics were studied such as leaf area dry weight ratio (LADWR), Dust retaining capacity (DRC) and Chlorophyll content. In the study the effects of marble dust on selected tree species was observed. LADWR was recorded maximum 217.90 cm² g⁻¹ dry wt. in *Polyalthia longifolia* and minimum 98.74 cm² g⁻¹ dry wt. in *Ficus religiosa* in Nindar whereas The DRC was observed maximum 178.51 mg cm⁻² in *Azadirachta indica* as well as minimum recorded 66.41 mg cm⁻² in *Thevetia peruviana*. However the Total Chlorophyll Content was also determined and it was found maximum in *Bougainvillea* i.e. 2.949 mg g⁻¹(fresh wt.) whereas minimum 0.784 mg g⁻¹ (fresh wt.) in *Ficus religiosa* (Saini, et al., 2011).

5. Conclusion and Future Aspects

The physiological response of plants reveals many different direct routes of action through which dust can affect plants. Mining dust may also exacerbate secondary stresses, such as drought, insects and pathogens, or allow penetration of toxic metals or phytotoxic gaseous pollutants. Effect of mining dust on natural communities may alter the competitive balance between species in a community. These changes in the vegetation may also affect animal communities, from vertebrate's grazers to soil invertebrates. This may, for example, alter cycles of decomposition. Response of individual species may be positive or negative depending on the particular situation, and only detailed studies may reveal the main reason behind any observed changes. There have, unfortunately, been only a limited number of studies at the community level.

It is evident from this review of the literature that there are many gaps in our knowledge of the effects of mining dust. Until research into these areas is undertaken, the ability of those with responsibility for plant and crop protection, or preventing the deterioration of natural and semi-natural habitats, to address developments that may threaten sites will be inadequate. It is important therefore, that current trends in dust emissions are identified as well as the vegetation types that are likely to be affected by such emission.

Thus, it is evident from the several research reports that judicious use and presence of mining dust have toxic effect on plants, animals and other living organisms and affects the same after certain limits. Therefore, it is well needed to intensify the research program for better understanding of effect of mining dust on plants and allied areas to maintain the ecological harmony.

Reference

- [1] Asubiojo, O.I., Aina, P.O and Oluwole, A.F (1991), "Effect of cement production on the elemental composition of soil in the neighbourhood of tow cement factories", *Water Air Soil pollution*, 57-58, 819-828.
- [2] Bachman, G.R and Miller, W.B (1995), "Iron chelate inducible iron/manganese toxicity in zonal geranium", *J. Plant Nutri.*, 18, 1917-1929.
- [3] Clarimont, K.B, Hagar, W.G, Davis, E.A (1986), "Manganese toxicity to chlorophyll synthesis in tobacco callus", *Plant Physiolo.*, 80, 291-293.
- [4] Crawford, T.W, Stroehlein, J.L, Kuehl, R.O (1989), "Manganese and rates of growth and mineral accumulation in cucumber", *J. Am Soc. Horti. Sci.*, 114, 300-306.
- [5] Demirevska-Kepova K, Simova-Stoilova L, Stoyanova Z, Holzer R, Feller U (2004), "Biochemical changes in barley plants after excessive supply of copper and manganese", *Environ Exp. Bot.*, 52, 253-266.
- [6] Elamin, O.M and Wilcox, G.E (1986a), "Effect of Manganese nutrition on muskmelon growth and manganese toxicity", *J Am Soc Horti. Sci.*, 111, 582-587.
- [7] Elamin, O.M and Wilcox, G.E (1986b), "Effect of Manganese nutrition on watermelon growth and manganese toxicity". *J. Am. Soc. Horti. Sci.*, 111, 582-587.
- [8] Farmer, A.M (1993), "Effect of dust on vegetation- A review. Environmental pollution", 79, 63-75.
- [9] Fernandes, J.C, Henriques, F.S (1991), "Biochemical, physiological and structural effects of excess copper in plants", *Bot Rev*, 57, 247-273.
- [10] Fennelly, P.F. (1195), "Primary and secondary particulates as pollutants". *Journal Air pollut. Contr. Assoc.*, 25, 697-704.
- [11] Foy, C.D, Weil, R.R, Coradetti, C.A (1995), "Differential manganese tolerances of cotton genotypes in nutrient solution", *J. Plant Nutrition*, 18, 685-706.
- [12] Harris, P.R. (1975), "Quarrying in national parks". *Quarry management and products*. March 1975, 55-58.
- [13] Hegedus, A, Erdei, S, Horvath, G (2001), "Comparative studies of H₂O₂ detoxifying enzymes in green and greening barley seedlings under cadmium stress", *Plant Sci.*, 160, 1085-1093.
- [14] Horiguchi, T (1988), "Mechanism of manganese toxicity and tolerance of plants. IV. Effects of silicon on alleviation of manganese toxicity of rice plants", *Soil Sci. Plant Nutri.*, 34, 65-73.
- [15] Horst, J (1988), "Beschreibung der Gleichgewichtslage des ionenaustauschs an schwach sauren harzen mit hilfe eines models der oberflächenkomplexbildung", *Doctoral thesis, University of Karlsruhe, Kfk report*, 4464.
- [16] J. Chatterjee, and C. Chatterjee (2000), "Phytotoxicity of cobalt, chromium and copper in cauliflower", *Environmental Pollution*, vol. 109, no. 1, 69-74.
- [17] Kitao, M, Lei, T.T and Koike, T (1997), "Effect of manganese toxicity on photosynthesis of white birch (*Betula platyphylla* var. *japonica*) seedlings", *Physiol Plant*, 101, 249-256.
- [18] Le, Bot. J, Kirkby, E.A and Beusichem, M.L (1990), "Manganese toxicity in tomato plants: effects on cation uptake and distribution", *J. Plant Nutria.*, 13, 513-525.
- [19] Loneragan, J.F (1988), "Distribution and movement of manganese in plants". In: Graham RD, Hannam RJ, Uren NC (eds) *Manganese in soils and plants*, Kluwer, Dordrecht, 113-124.
- [20] Marschner, H (1995), "Mineral nutrition of higher plants", 2nd edn. *Academic press*, Toronto.
- [21] Mohnish Pichhode and Kumar Nikhil (2015) a, "Effect of Copper Dust on Photosynthesis Pigments Concentration in Plants Species", *International Journal of Engineering Research and Management (IJERM)*, Vol.2, No.2 pp.63-66, February, 2015 (Online).
- [22] Mohnish Pichhode and Kumar Nikhil (2015) b, "Effect of Copper Mining Dust on the Soil and vegetation in India: A Critical Review", *International Journal of Modern Sciences and Engineering Technology (IJMSET)*, Vol.2, No.2 pp.73-76, February, 2015 (Online).
- [23] Nriagu, J.O (1989), "A global assessment of natural sources of atmospheric trace metals". *Nature*, 338, 47-49.
- [24] Nikki Phipps (2000), "The Bulb-o-licious Garden" : A bulb lover's guide to growing your favorite plants. Ed. by Nikki Phipps (2000), Publisher: lulu.com (July 19, 2010), ISBN-10: 055755733X, ISBN-13: 978-0557557332.
- [25] Panda, A.K.S, Murlidhar, J and Shahoo, B.N (1996), "Seasonal water quality assessment of jajang iron and mining area", *Jr. of Industrial Pollution Control*, 12 (1), 9-14.
- [26] Peplow, D (1999), "Environmental impacts of mining in Eastern Washington. Center for water and watershed studies fact sheet", *University of Washington. Seattle*.
- [27] Prasad, M.S.V and Inamdar, J.A (1990), "Effect of cement kiln dust pollution on blackgram (*Vigna munga*, L. hepper)", *Proc. India Acad. Sci., (Plant Science)*, 100, 435-443.
- [28] Prasad, M.S.V, Subramanian, R.B and Inamdar, J.A (1991), "Effect of cement kiln dust on *Cajanus cajan* (L.) Mill. Species", *Indian Jr. Environ. Hlth*, 33, 11-21.
- [29] Reeves, R.D and Baker, A.J.M (2000), "Metal-accumulating plants", In: Raskin I, Ensley BD (eds) *Phytoremediation of toxic metals: using plants to clean up the environment*. Wiley, New York, 193-229.
- [30] Rohan, J. Lad and Jay, S. Samant (2012), "Studies on the Impact of Bauxite Mining Activities on Environment in Kolhapur District", *Proceeding of International Conference SWRDM-2012*, Department of Environmental Science, Shivaji University, Kolhapur, pp.188-192.
- [31] Saini Yashoda, Nagendra Bhardwaj, Radha Gautam, (2011), "Effect of marble dust on plants around Vishwakarma Industrial Area (VKIA) in Jaipur, India",

*J. Environ. Biol.*32, 209-212 (2011) ISSN: 0254-8704, CODEN: JEBIDP.

- [32] Saralabai VC (1993), "Effect of cement kiln exhaust (Electrostatic precipitator dust) on growth, root nodule biochemistry and crop productivity in legumes through simulation studies" *Ph.D thesis, Bharathidasan University, Thiruchirapalli.* 1993.
- [33] Stanton, W. (1989), "Bleak prospects for limestone", *New Scientist*, 122(1664), 56--60.
- [34] Stadtman, E.R and Oliver, C.N (1991), "Metal-catalyzed oxidation of proteins", *J. Bio. Chem.* 266: 2005–2008.
- [35] Stern, A.C (1976), "Air pollution, Measurement, monitoring and surveillance of air pollution", 3rd ed., *Academic press, New York.*
- [36] Thomas, F, Malick, C, Endreszl, E.C, Davies, K.S (1998), "Distinct response to copper stress in the halophyte, *Mesembryanthemum crystallium*", *Physiol. Plant*, 102, 360-368.
- [37] Wu, S (1994), "Effect of manganese excess on the soybean plant cultivated under various growth conditions", *J. Plant Nutri.*, 17, 993-1003.

Author Profile



Jyotish Katare, M.Sc., Department of Biotechnology, Sardar Patel College of Technology, Balaghat affiliated to Rani Durgavati Vishwavidyalaya, Jabalpur, M.P. done his project work at EMG, CSIR-CIMFR, Dhanbad, Jharkhand, India in the year 2013 and publishing his experimental finding which is very significant and important work. This review paper will helps in understanding the effects of copper and other mining present at district Balaghat, M.P through dust showing effects on plants and soil. He further grown plants on the copper overburden dump top material as growing media with or without soil and Cowdung manure in different combination shown significant cumulative effects on the plant growth which has already been published.



Mohnish Pichhode, M.Sc. in Biotechnology, Department of Biotechnology, Sardar Patel College of Technology, Balaghat affiliated to Rani Durgavati Vishwavidyalaya, Jabalpur, M.P done his project work at EMG, CSIR-CIMFR, Dhanbad, Jharkhand, India in the year 2013 and published very significant and important review paper as well as an experimental work in the second paper which had indeed helped in understanding the effects of copper mining in nearby vegetation and soil within the district Balaghat, M.P.



Dr. Kumar Nikhil, Ph.D in Env.Sc. & Engg., Principal Scientist at Environmental Management Group, CSIR-Central Institute of Mining and Fuel Research (CIMFR), Barwa Road, Dhanbad - 826 015, Jharkhand, India has gained more than 30 years of research experience involved in more than 60 projects in different capacity. More than 135 scientific publications on his name. Guided more than 65 students of B.Sc, M.Sc, B.Tech & M.Tech, Ph.D students in their project and research works.